

Measurements of CKM angle ϕ_1 with charmless penguins at Belle

J. Dalseno for the Belle Collaboration

High Energy Accelerator Research Organization (KEK), Tsukuba, JAPAN

We present measurements of time-dependent CP violation parameters in $B^0 \rightarrow K^0\pi^0$, $B^0 \rightarrow K_S^0\pi^+\pi^-$ and $B^0 \rightarrow K_S^0K^+K^-$ decays. The latter two are extracted using time-dependent Dalitz plot analyses. These results are obtained from a large data sample that contains 657×10^6 $B\bar{B}$ pairs collected at the $\Upsilon(4S)$ resonance with the Belle detector at the KEKB asymmetric-energy e^+e^- collider.

1. INTRODUCTION

CP violation in the Standard Model (SM) arises from an irreducible complex phase in the Cabibbo-Kobayashi-Maskawa (CKM) quark-mixing matrix [1, 2]. Of recent interest is CP violation in $b \rightarrow q\bar{q}s$ transitions which proceeds by loop diagrams that may be affected by new particles in various extensions of the SM. Furthermore, the CP asymmetries in $b \rightarrow q\bar{q}s$ transitions are predicted in the SM to be slightly higher than those observed in $b \rightarrow c\bar{c}s$ transitions. However, current experimental measurements [3] tend to be lower than those for $b \rightarrow c\bar{c}s$ transitions motivating more precise experimental determinations.

The decay of the $\Upsilon(4S)$ produces a $B\bar{B}$ pair of which one (B_{Rec}^0) may be fully reconstructed while the other (B_{Tag}^0) may reveal its flavour. The proper time interval between B_{Rec}^0 and B_{Tag}^0 is defined as $\Delta t \equiv t_{\text{Rec}} - t_{\text{Tag}}$ and from coherent $B\bar{B}$ production in the $\Upsilon(4S)$ decay, the time-dependent decay rate for a quasi-two-body mode when B_{Tag}^0 possesses flavour q (B^0 : $q = +1$, $\bar{B}^0$: $q = -1$), is given by [4]

$$|A(\Delta t, q)|^2 = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} \left[1 + q(\mathcal{A}_{CP} \cos \Delta m_d \Delta t + \mathcal{S}_{CP} \sin \Delta m_d \Delta t) \right], \quad (1)$$

where τ_{B^0} is the B^0 lifetime and Δm_d is the $B^0\bar{B}^0$ mass difference. This assumes no CP violation in mixing, $|q/p| = 1$, and that the $B^0\bar{B}^0$ lifetime difference is negligible. The parameter, $\mathcal{A}_{CP}$, denotes the direct CP violating component and $\mathcal{S}_{CP}$ represents mixing-induced CP violation. For time-dependent Dalitz plot analyses, the time-dependent decay rate is written as

$$|A(\Delta t, q)|^2 = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} \left[(|A|^2 + |\bar{A}|^2) - q(|A|^2 - |\bar{A}|^2) \cos \Delta m_d \Delta t + 2q\Im(\bar{A}A^*) \sin \Delta m_d \Delta t \right]. \quad (2)$$

The Dalitz-dependent amplitudes, A , can be written in the isobar approximation as a superposition of intermediate decay channels, i ,

$$A(s_+, s_-) = \sum_i a_i F_i(s_+, s_-), \quad \bar{A}(s_-, s_+) = \sum_i \bar{a}_i \bar{F}_i(s_-, s_+), \quad (3)$$

where $a_i \equiv a_i(1 + c_i)e^{i(b_i + d_i)}$ for A and $\bar{a}_i \equiv a_i(1 - c_i)e^{i(b_i - d_i)}$ for $\bar{A}$ are complex coefficients describing the relative magnitudes and phases between the decay channels. The form factors, $F_i(s_+, s_-)$, depend on the Dalitz plot coordinates, $s_{\pm}$, and describe the invariant mass and angular distribution probabilities. For a CP eigenstate, i , the time-dependent CP violation parameters can be calculated as

$$\mathcal{A}_{CP}(i) \equiv \frac{|\bar{a}_i|^2 - |a_i|^2}{|\bar{a}_i|^2 + |a_i|^2} = \frac{-2c_i}{1 + c_i^2}, \quad \text{and} \quad -\eta_i \mathcal{S}_{CP}(i) \equiv \frac{-2\Im(\bar{a}_i a_i^*)}{|a_i|^2 + |\bar{a}_i|^2} = \frac{1 - c_i^2}{1 + c_i^2} \sin 2\phi_1^{\text{eff}}(i) \quad (4)$$

and $\phi_1^{\text{eff}}(i) \equiv \arg(a_i \bar{a}_i^*)/2 = d_i$ is directly accessible as a fit parameter. Consequently, both $\mathcal{A}_{CP}(i)$ and $\mathcal{S}_{CP}(i)$ are restricted to reside in the physical region. The relative fraction of each component can be calculated with,

$$f_i = \frac{(|a_i|^2 + |\bar{a}_i|^2) \int F_i(s_+, s_-) F_i^*(s_+, s_-) ds_+ ds_-}{\int (|A|^2 + |\bar{A}|^2) ds_+ ds_-}. \quad (5)$$

2. DATASET, DETECTOR AND BASIC ANALYSIS TECHNIQUE

These measurements of CP violating parameters are based on 657×10^6 $B\bar{B}$ pairs collected with the Belle detector [5] at the KEKB asymmetric-energy e^+e^- (3.5 on 8 GeV) collider [6]. Operating with a peak luminosity that exceeds $1.7 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$, the collider produces the $\Upsilon(4S)$ resonance ($\sqrt{s} = 10.58 \text{ GeV}$) with a Lorentz boost of $\beta\gamma = 0.425$, opposite to the positron beam direction, z , which usually decays into a $B\bar{B}$ pair.

Reconstructed B candidates are described with two kinematic variables: the beam-constrained mass, $M_{bc} \equiv \sqrt{(E_{\text{beam}}^{\text{CMS}})^2 - (p_B^{\text{CMS}})^2}$ and the energy difference, $\Delta E \equiv E_B^{\text{CMS}} - E_{\text{beam}}^{\text{CMS}}$ where $E_{\text{beam}}^{\text{CMS}}$ is the beam energy and E_B^{CMS} (p_B^{CMS}) is the energy (momentum) of the B meson all evaluated in the centre-of-mass system (CMS). The dominant background in the reconstruction of B_{Rec}^0 is from continuum ($e^+e^- \rightarrow q\bar{q}$) events. Since their topology tends to be jet-like in contrast to the spherical $B\bar{B}$ decay, continuum can be suppressed with a Fisher discriminant based on modified Fox-Wolfram moments [7]. This discriminant is combined with the polar angle of the B candidate in the CMS to form a likelihood ratio, $\mathcal{R}_{S/B}$, which separates continuum from $B\bar{B}$ events.

Since the B^0 and $\bar{B}^0$ mesons are approximately at rest in the $\Upsilon(4S)$ Enter-of-Mass System (CMS), the difference in decay time between the $B\bar{B}$ pair, Δt , can be determined from the displacement in z between the final state decay vertices, $\Delta t \simeq (z_{\text{Rec}} - z_{\text{Tag}})/\beta\gamma c \equiv \Delta z/\beta\gamma c$. To obtain the Δt distribution, we reconstruct the tag-side vertex from the tracks not used to reconstruct B_{Rec}^0 [8] and employ the flavour tagging routine described in Ref. [9]. The tagging information is represented by two parameters, the B_{Tag}^0 flavour, q and r . The parameter, r , is an event-by-event, MC determined flavour-tagging dilution factor that ranges from $r = 0$ for no flavour discrimination to $r = 1$ for unambiguous flavour assignment.

3. TIME-DEPENDENT CP VIOLATION MEASUREMENT IN $B^0 \rightarrow K^0\pi^0$

Direct CP violation has been observed in $B^0 \rightarrow K^+\pi^-$ [10] and is found to be significantly different from that in $B^\pm \rightarrow K^\pm\pi^0$. This unexpected result may indicate the presence of new physics (NP) or poor understanding of strong interaction effects in B decays. A model-independent test for NP is possible via an isospin sum rule with high statistics [11] which gives a relation among the direct CP violation asymmetries measured in all possible $B \rightarrow K\pi$ modes, i.e. $B^0 \rightarrow K^0\pi^0$, $K^+\pi^-$, $B^+ \rightarrow K^+\pi^0$ and $K^0\pi^+$,

$$\mathcal{A}_{K^+\pi^-} + \mathcal{A}_{K^0\pi^+} \frac{\mathcal{B}(K^0\pi^+)\tau_{B^0}}{\mathcal{B}(K^+\pi^-)\tau_{B^+}} = \mathcal{A}_{K^+\pi^0} \frac{2\mathcal{B}(K^+\pi^0)\tau_{B^0}}{\mathcal{B}(K^+\pi^-)\tau_{B^+}} + \mathcal{A}_{K^0\pi^0} \frac{2\mathcal{B}(K^0\pi^0)}{\mathcal{B}(K^+\pi^-)}. \quad (6)$$

Here, $\mathcal{B}$ represents the branching ratio of each decay mode and τ_{B^+} is the lifetime of the charged B meson. The sum rule's theoretical precision is determined by SU(2) flavour symmetry, i.e. a few %, therefore the sum rule provides a clean test for new physics. As a fundamental input, the CP violation measurement in $B^0 \rightarrow K^0\pi^0$ is currently the least known, experimentally. Since the branching fractions and CP asymmetries of other $B \rightarrow K\pi$ decay modes have been measured to good precision [12], $\mathcal{A}_{K^0\pi^0}$ is predicted with a small error. In addition to the $B^0 \rightarrow K_S^0\pi^0$ mode, we measure CP asymmetry in $B^0 \rightarrow K_L^0\pi^0$ decay for the first time, in order to maximize sensitivity to the direct CP violation parameter, $\mathcal{A}_{K^0\pi^0}$.

The signal yield of $B^0 \rightarrow K_S^0\pi^0$ events is found from a three-dimensional extended unbinned maximum likelihood fit to M_{bc} , ΔE and $\mathcal{R}_{S/B}$ to be 657 ± 37 events. For $B^0 \rightarrow K_L^0\pi^0$, M_{bc} can be calculated from the direction of the K_L^0 cluster while ΔE cannot be calculated. The signal yield of 285 ± 52 events is extracted from a two-dimensional fit to M_{bc} and $\mathcal{R}_{S/B}$ and has a significance of 3.7σ including systematic uncertainties.

As the vertex position of $B^0 \rightarrow K_L^0\pi^0$ cannot be determined and the vertex reconstruction efficiency of $B^0 \rightarrow K_S^0\pi^0$ is $\sim 33\%$ due to the long lifetime of the K_S^0 , these events can still be used to calculate $\mathcal{A}_{CP}$ by integrating Eq. 1 over Δt . We extract the CP parameters,

$$\begin{aligned} \mathcal{A}_{CP} &= +0.14 \pm 0.13 \text{ (stat)} \pm 0.06 \text{ (syst)}, \\ \mathcal{S}_{CP} &= +0.67 \pm 0.31 \text{ (stat)} \pm 0.08 \text{ (syst)} \end{aligned} \quad (7)$$




Figure 1: The left plot shows the background subtracted fit to Δt for the good tags region, $0.5 < r \leq 1.0$, where the solid (dashed) curve represents B^0 ($\bar{B}^0$) tags. The right plot shows the $B^0 \bar{B}^0$ raw asymmetry, $(N_{B^0} - N_{\bar{B}^0}) / (N_{B^0} + N_{\bar{B}^0})$, where N_{B^0} ($N_{\bar{B}^0}$) is the number of signal B^0 ($\bar{B}^0$) tags in Δt .

and the fit results for the Δt component is shown in Fig. 1. We find that the mixing induced component is consistent with charmonium, $\mathcal{S}_{CP}(b \rightarrow c\bar{c}s) = 0.681 \pm 0.025$ [3], and that there is a 1.9σ deviation between our measurement of $\mathcal{A}_{CP}$ and the expectation from the isospin sum rule, $\mathcal{A}_{CP} = -0.17 \pm 0.06$ [11].

4. TIME-DEPENDENT DALITZ PLOT CP VIOLATION MEASUREMENT IN $B^0 \rightarrow K_S^0 \pi^+ \pi^-$

The signal yield of $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ events is found from a one-dimensional fit to ΔE to be 1944 ± 98 events and the resonances considered in the signal model (Eq. 3) are the $K^{*\pm}(892)$, $K_0^{*\pm}(1430)$, $\rho^0(770)$, $f_0(980)$, $f_2(1270)$, $f_X(1300)$ and a non-resonant component. We obtain the CP parameters,

Solution 1: $-2 \log \mathcal{L} = 18472.5$

$$f(K_0^{*+}(1430)\pi^-) = 61.7 \pm 10.4\%$$

$$\mathcal{A}_{CP}(\rho^0(770)K_S^0) = +0.03_{-0.24}^{+0.23} \pm 0.11 \pm 0.10,$$

$$\phi_1^{\text{eff}}(\rho^0(770)K_S^0) = (+20.0_{-8.5}^{+8.6} \pm 3.2 \pm 3.5)^\circ,$$

$$\mathcal{A}_{CP}(f_0(980)K_S^0) = -0.06 \pm 0.17 \pm 0.07 \pm 0.09,$$

$$\phi_1^{\text{eff}}(f_0(980)K_S^0) = (+12.7_{-6.5}^{+6.9} \pm 2.8 \pm 3.3)^\circ,$$

Solution 2: $-2 \log \mathcal{L} = 18465.0$

$$f(K_0^{*+}(1430)\pi^-) = 17.4 \pm 5.0\%$$

$$\mathcal{A}_{CP}(\rho^0(770)K_S^0) = -0.16 \pm 0.24 \pm 0.12 \pm 0.10,$$

$$\phi_1^{\text{eff}}(\rho^0(770)K_S^0) = (+22.8 \pm 7.5 \pm 3.3 \pm 3.5)^\circ,$$

$$\mathcal{A}_{CP}(f_0(980)K_S^0) = +0.00 \pm 0.17 \pm 0.06 \pm 0.09,$$

$$\phi_1^{\text{eff}}(f_0(980)K_S^0) = (+14.8_{-6.7}^{+7.3} \pm 2.7 \pm 3.3)^\circ, \quad (8)$$

where the first error is statistical, the second is systematic and the third is the Dalitz plot signal model uncertainty. Figure 2 shows the fit results. The high $K_0^{*+}(1430)\pi^-$ fraction of Solution 1 is in agreement with some phenomenological estimates [13] and may also be favoured by the total $K - \pi$ S -wave phase shift when compared with that measured by LASS [14]. As the likelihood difference is not found to be significant, we do not rule between these solutions where $\mathcal{A}_{CP}$ is consistent with null asymmetry and ϕ_1^{eff} agrees with charmonium.

5. TIME-DEPENDENT DALITZ PLOT CP VIOLATION MEASUREMENT IN $B^0 \rightarrow K_S^0 K^+ K^-$

The signal yield of $B^0 \rightarrow K_S^0 K^+ K^-$ is extracted to be 1269 ± 51 events from a two-dimensional fit to M_{bc} and ΔE in each r -bin. We consider the $f_0(980)$, $\phi(1020)$, $f_X(1500)$, χ_{c0} , and a non-resonant component in the signal model. Four solutions with similar likelihood are found where two solutions arise from the interference between $f_0(980)$ and the non-resonant component and the other two from $f_X(1500)$ and the non-resonant component. Using external information from $B^0 \rightarrow K_S^0 \pi^+ \pi^-$, if the $f_X(1500)$ is the $f_0(1500)$ for both $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ and $B^0 \rightarrow K_S^0 K^+ K^-$, the ratio of branching fractions, $\mathcal{B}(f_0(1500) \rightarrow \pi^+ \pi^-) / \mathcal{B}(f_0(1500) \rightarrow K^+ K^-)$, prefers the solution with the low $f_X(1500)K_S^0$ fraction. Similarly the ratio, $\mathcal{B}(f_0(980) \rightarrow \pi^+ \pi^-) / (\mathcal{B}(f_0(980) \rightarrow \pi^+ \pi^-) + \mathcal{B}(f_0(980) \rightarrow K^+ K^-))$




Figure 2: Time-dependent Dalitz plot fit results for $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ in (a), the $\rho^0(770)$ region and (b), the $f_0(980)$ region. The top plots show the Δt distribution for B^0 (solid) and $\bar{B}^0$ (dashed) tags. These plots contain only good tags, $0.5 < r \leq 1.0$ and the dotted curve represents the background contribution. The bottom plots show the $B^0 \bar{B}^0$ raw asymmetry, $(N_{B^0} - N_{\bar{B}^0}) / (N_{B^0} + N_{\bar{B}^0})$, where N_{B^0} ($N_{\bar{B}^0}$) is the number of B^0 ($\bar{B}^0$) tags in Δt .

prefers the solution with the low $f_0(980)K_S^0$ fraction. The preferred solution is,

$$\begin{aligned}
 \mathcal{A}_{CP}(f_0(980)K_S^0) &= -0.02 \pm 0.34 \pm 0.08 \pm 0.09 \\
 \phi_1^{\text{eff}}(f_0(980)K_S^0) &= (28.2_{-9.9}^{+9.8} \pm 2.0 \pm 2.0)^\circ \\
 \mathcal{A}_{CP}(\phi(1020)K_S^0) &= +0.31_{-0.23}^{+0.21} \pm 0.04 \pm 0.09 \\
 \phi_1^{\text{eff}}(\phi(1020)K_S^0) &= (21.2_{-10.4}^{+9.8} \pm 2.0 \pm 2.0)^\circ
 \end{aligned} \tag{9}$$

where the first error is statistical, the second is systematic and the third is the Dalitz plot signal model uncertainty. Figure 3 shows the fit results where $\mathcal{A}_{CP}$ is consistent with null asymmetry and ϕ_1^{eff} agrees with charmonium.

Acknowledgments

We thank the KEKB group for excellent operation of the accelerator, the KEK cryogenics group for efficient solenoid operations, and the KEK computer group and the NII for valuable computing and SINET3 network support. We acknowledge support from MEXT and JSPS (Japan); ARC and DEST (Australia); NSFC (China); DST (India); MOEHRD and KOSEF (Korea); KBN (Poland); MES and RFAAE (Russia); ARRS (Slovenia); SNSF (Switzerland); NSC and MOE (Taiwan); and DOE (USA).

References

- [1] N. Cabibbo, Phys. Rev. Lett. **8**, 214 (1964).
- [2] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [3] E. Barberio *et al.* (Heavy Flavor Averaging Group), arXiv:0808.1297 [hep-ex] and online update for Winter 2008 at <http://www.slac.stanford.edu/xorg/hfag>.




Figure 3: Time-dependent Dalitz plot fit results for $B^0 \rightarrow K_S^0 K^+ K^-$ in the $\phi(1020)$ region. The left plot show the Δt distribution for B^0 (blue) and $\bar{B}^0$ (red) tags for good tags, $0.5 < r \leq 1.0$. The right plot shows the $B^0 \bar{B}^0$ raw asymmetry

- [4] A. B. Carter and A. I. Sanda, Phys. Rev. Lett. **45**, 952 (1980); A. B. Carter and A. I. Sanda, Phys. Rev. D **23**, 1567 (1981); I. I. Bigi and A. I. Sanda, Nucl. Phys. **193**, 85 (1981).
- [5] A. Abashian *et al.* (Belle Collab.), Nucl. Instr. and Meth. A **479**, 117 (2002).
- [6] S. Kurokawa and E. Kikutani, Nucl. Instr. and Meth. A **499**, 1 (2003), and other papers included in this volume.
- [7] K. Abe *et al.* (Belle Collaboration), Phys. Rev. Lett. **87**, 101801 (2001); K. Abe *et al.* (Belle Collaboration), Phys. Lett. **B 511**, 151 (2001); S. H. Lee, K. Suzuki, *et al.* (Belle Collaboration), Phys. Rev. Lett. **91**, 261801 (2003).
- [8] H. Tajima *et al.* Nucl. Instr. and Meth. A **533**, 370 (2004).
- [9] H. Kakuno *et al.*, Nucl. Instr. and Meth. A **533**, 516 (2004).
- [10] B. Aubert *et al.* (BaBar Collab.), Phys. Rev. Lett. **99**, 021603 (2007); S.-W. Lin *et al.* (Belle Collab.), Nature **452**, 332 (2008).
- [11] M. Gronau, Phys. Lett. B **627**, 82 (2005).
- [12] C. Amsler (Particle Data Group), Phys. Lett. B **567**, 1 (2008).
- [13] V. L. Chernyak, Phys. Lett. B **509**, 273 (2001).
- [14] D. Aston *et al.* (LASS Collab.), Nucl. Phys. B **296**, 493 (1988).